

# COLONOSCOPY - GOLYTELY PREP INSTRUCTIONS

## **Patient Responsibility**

Please check with your insurance company to fulfill any precertification requirements that you may have. Also, please be aware that there are some circumstances that may change the type of colonoscopy that is billed to your insurance. If a biopsy is done or a polyp is removed, this may change a screening colonoscopy to a diagnostic colonoscopy. This could result in higher out-of-pocket expenses. Please contact your insurance company for additional information.

Procedure	Date:	
-----------	-------	--

## **PURCHASE AT THE PHARMACY:**

1. Prescription for Golytely

## **MEDICATIONS:**

If you are taking prescription medications (including anticoagulant/antiplatelet), please discuss with your primary care physician which medications, if any, need to be held prior and/or taken the morning of the procedure.

Medication Name - NSAID Medication	HOLD	RESTART
Advil/Motrin (Ibuprofen)	Hold 5-7 days if able	Day after procedure
Aleve (Naproxen)	Hold 5-7 days if able	Day after procedure
Celebrex (Celecoxib)	Hold 5-7 days if able	Day after procedure
Diclofenac (Voltaren)	Hold 5-7 days if able	Day after procedure
Excedrin	Hold 5-7 days if able	Day after procedure
Meloxicam (Mobic)	Hold 5-7 days if able	Day after procedure
Toradol (Ketoralac)	Hold 5-7 days if able	Day after procedure
Aspirin (325 mg or higher)	Hold day of procedure	Day after procedure
A :: (I I :: ) O1	Hold day of propodure	Day after procedure
Aspirin (baby aspirin) - 81 mg	Hold day of procedure	Day after procedure
Medication Name – Anticoagulants	Recommendations ONLY - patient needs t	to confirm hold with Prescribing I
Medication Name - Anticoagulants Eliquis (Apixaban)	Recommendations ONLY - patient needs t	to confirm hold with Prescribing I
Medication Name - Anticoagulants Eliquis (Apixaban)	Recommendations ONLY - patient needs t	to confirm hold with Prescribing I
Medication Name – Anticoagulants	Recommendations ONLY - patient needs t	to confirm hold with Prescribing I
Medication Name - Anticoagulants Eliquis (Apixaban) Lovenox (Enoxaparin)	Recommendations ONLY - patient needs to Hold 48 hours Hold 48 hours	to confirm hold with Prescribing I 48-72 hours 48-72 hours 48-72 hours
Medication Name - Anticoagulants Eliquis (Apixaban) Lovenox (Enoxaparin) Pradaxa (Dabigatran)	Recommendations ONLY - patient needs to Hold 48 hours Hold 48 hours Hold 48 hours	to confirm hold with Prescribing I 48-72 hours 48-72 hours 48-72 hours
Medication Name - Anticoagulants  Eliquis (Apixaban)  Lovenox (Enoxaparin)  Pradaxa (Dabigatran)  Warfarin (Coumadin)	Recommendations ONLY - patient needs to Hold 48 hours Hold 48 hours Hold 48 hours Hold 5 days (confirm INR = 1.5 days prior)</td <td>48-72 hours 48-72 hours 48-72 hours Restart evening of procedure Confirm with Prescribing MD</td>	48-72 hours 48-72 hours 48-72 hours Restart evening of procedure Confirm with Prescribing MD
Medication Name - Anticoagulants  Eliquis (Apixaban)  Lovenox (Enoxaparin)  Pradaxa (Dabigatran)  Warfarin (Coumadin)  Xarelto (Rivaroxaban)  Medication Name -	Recommendations ONLY - patient needs to Hold 48 hours Hold 48 hours Hold 48 hours Hold 5 days (confirm INR = 1.5 days prior) Confirm with Prescribing MD</td <td>48-72 hours 48-72 hours 48-72 hours Restart evening of procedure Confirm with Prescribing MD</td>	48-72 hours 48-72 hours 48-72 hours Restart evening of procedure Confirm with Prescribing MD

# COLONOSCOPY PREPARATION INSTRUCTIONS (continued)

### MEDICATIONS continued...

## **Diabetic Patients:**

Please do not take your morning dose of oral diabetic medication the day of the procedure. If you are on insulin, please take only one-half (1/2) of your usual morning insulin dose on the day of the procedure.

#### **SGLT-2 Medications for Diabetic Patients:**

It is very important that you adhere to the medication hold for SGLT-2 medications, due to euglycemic diabetic ketoacidosis.

For patients taking any of the following SGLT-2 medications – Empagliflozin (Jardiance), Canagliflozin (Invokana), Dapagliflozin (Farxiga), Bexagliflozin (Brenzavvy), Sotagliflozin (Inpefa)

• Hold medication for 3 days prior to surgery.

Ertugliflozin (Steglatro)

• Hold medication for 4 days prior to surgery.

## **GLP1 Medications for Diabetic/Weight Loss Patients:**

It is very important that you adhere to the medication holds for GLP1 medications. Due to the delay in gastric emptying, these medications can increase the risk of regurgitation and pulmonary aspiration during anesthesia and moderate sedation.

For patients taking any of the following GLP1 medications - Dulaglutide (Trulicity- Weekly), Exenatide extended release (Bydureon bcise- Weekly), Exenatide (Byetta- Twice Daily), Liraglutide (Victoza and Saxenda- daily), Lixisenatide (Adlyzin- Daily), Semaglutide (Ozempic and Wegovy- Weekly), Tirzepatide (Mounjaro- Weekly), Semaglutide (Rybelsus-Taken by mouth once daily)

- If you are taking a daily dosing of a GLP1 (usually orals), hold the medication the morning of the procedure and resume your regular schedule following the procedure.
- If you are taking a weekly dosing of a GLP1 (usually injectables), hold the medication for one week prior to the
  procedure and resume the regular schedule following the procedure.

**GLP1 Medication Dietary Instructions -** All patients who are receiving anesthesia and are on a GLP1 should be on a clear liquid diet for 24 hours prior to surgery. You may continue drinking clear liquids up until 4 hours before your procedure. Clear liquids include black coffee or tea, clear fruit juices (apple or white grape), broth or bouillon, non-red Jell-O, or essentially any fluid you can see through.

### TWO DAYS BEFORE THE PROCEDURE:

Eat a low fiber diet. No fruits or vegetables, no whole wheat foods, no nuts, seeds or popcorn. No fiber supplements. Drink at least eight glasses of water during the day.

## DAY BEFORE THE PROCEDURE:

- 1. No solid food or milk products are allowed. You may have clear liquids: examples include crystal light, 7-UP, Gatorade, Lemonade, tea, coffee, soft drinks, clear/strained fruit drinks, flavored drinks, Kool-Aid, strained broth (canned or bouillon cubes) and Jell-O. Do not eat anything red in color.
- 2. You may continue drinking clear liquids up until 4 hours before your procedure.
- 3. Mix the Golytely with warm tap water to the fill line and refrigerate in the morning. You may add crystal light drink mix (not red in color) to improve the taste. Lemonade crystal light works best.
- **4.** Begin at 2:00 PM by drinking one 8-ounce glass of Golytely about every ten minutes until the container is consumed. You will begin going to the bathroom about one hour after beginning the prep.
- 5. Additional helpful instructions: wear warm clothing while taking the prep as you may feel chilled, drink the prep through a straw to help with the taste, chew gum during the prep to help with the taste and diminish nausea sensation.

The bowel prep is very important. If you do not completely empty your colon, this will limit the quality of your examination and may potentially mean taking an additional prep and returning for another examination sooner than would otherwise be required.

If you have problems with constipation on a daily basis, the bowel prep may not work as well so we encourage you to start your bowel prep as soon as possible the day before the procedure. If you have not started having bowel movements by early evening, please call so additional instructions can be given.

## COLONOSCOPY PREPARATION INSTRUCTIONS (continued)

## **DAY OF PROCEDURE:**

- 1. **Driver Information:** If you require anesthesia or sedation for your procedure, you must have a family member or friend bring you home from the surgery center. Public transportation, including Uber or Lyft type service, is not acceptable unless you are accompanied by a responsible adult who will be caring for you.
- 2. Nothing to eat or drink 4 hours prior to your procedure.
- 3. You will be given medication during the procedure that will make you sleepy.
- 4. The procedure takes about 30 minutes, but you will remain at the facility for about one hour after the procedure.
- 5. The most serious risks of the procedure are bleeding and perforation of the intestine. These are rare, but can be serious, and may require surgery to fix.
- 6. Please bring a Photo ID.

**RESULTS:** All finalized results are released to MyChart instantly. You may have access to your results before the provider is able to review them. Your provider will review the results and provide commentary within 1 week of the results finalizing. Our office will contact you via MyChart message, phone call, or letter regarding the results and next steps.

#### HOSPITAL STANDARD OF PRACTICE

Do not smoke after noon the day before your procedure. Remove all makeup, including mascara and nail polish, before coming to the hospital. Dentures, partials, contact lenses, hearing aids and piercings may be removed prior to your procedure. Please leave all valuables and jewelry at home. You are required to have a responsible adult drive you home. Bring a list of your current medications and supplements to the hospital with you, as well as your insurance cards and photo id. If you are an insulin dependent diabetic, your physician may choose to make your appointment early in the day. If you develop a cold, fever, sore throat, or any other illness, please notify our office before you come to the hospital. Your physician will decide if your procedure should be rescheduled. Firearms and weapons are prohibited on the hospital premises except in your locked motor vehicle.

If you have any questions or concerns, please do not hesitate to call our office at (920) 731-8131 or 1-800-574-3872.

## FOX VALLEY SURGICAL SPECIALISTS COLONOSCOPY INFORMATION

Fox Valley Surgical Specialists has created this information to help you sort through some of the confusion that can be associated with billing your insurance for colonoscopy procedures.

## Here are some helpful things to remember:

- There are some circumstances that may change the type of colonoscopy that is billed to your insurance company.
   If a biopsy is done or a polyp removed, this may change a <u>screening</u> colonoscopy to a <u>diagnostic</u> colonoscopy.
   This could result in higher out-of-pocket expenses. Please contact your insurance company for additional information about your individual policy and how it will process each type of colonoscopy.
- 2. Fox Valley Surgical Specialists <u>does not</u> contact your insurance company prior to your procedure. We ask each patient to be responsible for knowing his/her insurance company's guidelines and following through with any precertification requirements.
- 3. You will receive separate bills from your doctor's office and the facility where you had your procedure done. You may also receive a bill from the laboratory, pathology and/or anesthesiology. Fox Valley Surgical Specialists can only provide you with information associated with our fees.
- 4. Your doctor cannot add, delete or change a diagnosis (billing code) to facilitate better insurance coverage.
- 5. We are contractually and legally required to report any and all of our findings.
- 6. When your doctor dictates his operative note, it is a binding, legal document that cannot be altered for the sole purpose of coverage determination. We ask each patient to understand and respect our responsibility to report his/her procedure completely and honestly to the insurance company.

If you do not have insurance coverage for your procedure, please call to discuss our fees and payment arrangements. This is important to do prior to your colonoscopy to avoid any unexpected financial issues. If you have any additional questions regarding this information, please contact The Fox Valley Surgical Specialists Business Office at: 920-731-8289, Monday-Thursday, from 7:00 am to 5:00 pm and Friday, from 7:00 am to 4:00 pm.

Thank you for choosing Fox Valley Surgical Specialists for your healthcare needs.